

Deep infrared imaging and spectroscopy of the nearby late M-dwarf DENIS-P J104814–395606^{*}

R. NEUHÄUSER¹, E.W. GUENTHER², J. ALVES³, N. GROSSO¹, CH. LEINERT⁴, TH. RATZKA⁴, TH. OTT¹, M. MUGRAUER¹, F. COMERÓN³, A. ECKART⁵ AND W. BRANDNER³

¹ MPI für extraterrestrische Physik, D-85740 Garching, Germany

² Thüringer Landessternwarte Tautenburg, Sternwarte 5, D-07778 Tautenburg, Germany

³ European Southern Observatory, Karl-Schwarzschild-Straße 2, D-85748 Garching, Germany

⁴ MPI für Astronomie, Königstuhl, D-69117 Heidelberg, Germany

⁵ I. Physikalisches Institut, Universität zu Köln, Germany

Received date will be inserted by the editor; accepted date will be inserted by the editor

Abstract. We obtained deep H- and K-band images of DENIS-P J104814–395606 using SofI and the speckle camera SHARP-I at the ESO-3.5m-NTT as well as QUIRC at the Mauna Kea 2.2m telescope between December 2000 and June 2001. The target was recently discovered as nearby M9-dwarf among DENIS sources (Delfosse et al. 2001). We detect parallactic motion on our images and determine the distance to be 4.6 ± 0.3 pc, more precise than previously known. From the available colors, the distance, and the spectral type, we conclude from theoretical models that the star has a mass of ~ 0.075 to $0.09 M_{\odot}$ and an age of ~ 1 to 2 Gyrs. We also obtained H- and K-band spectra of this star with ISAAC at the VLT. A faint companion candidate is detected $6''$ NNW of the star, which is 6.4 ± 0.5 mag fainter in H. However, according to another image taken several month later, the companion candidate is not co-moving with the M9-dwarf. Instead, it is a non-moving background object. Limits for undetected companion candidates are such that we can exclude any stellar companions outside of $\sim 0.25''$ (1 AU), any brown dwarf companions (above the deuterium burning mass limit) outside of $\sim 2''$ (9 AU), and also any companion down to $\sim 40 M_{\text{Jup}}$ with $\geq 0.15''$ (0.7 AU) separation, all calculated for an age of 2 Gyrs. Our observations show that direct imaging of sub-stellar companions near the deuterium burning mass limit in orbit around nearby ultra-cool dwarfs is possible, even with separations that are smaller than the semi-major axis of the outermost planet in our solar system, namely a few tens of AU.

Key words: Editorial notes – instruction for authors

1. Introduction

The object DENIS-P J104814–395606 (hereafter Denis 1048–39) was discovered as very red source in the DEep Near-Infrared Survey (DENIS, Epchtein 1997) by Delfosse et al. (2001): $I=12.67 \pm 0.05$, $J=9.59 \pm 0.05$, and $K_s = 8.58 \pm 0.05$ mag. Together with its USNO B- and R-band colors ($B=19.0 \pm 0.3$ and $R=15.7 \pm 0.2$ mag), it was classified as late M-type star; a follow-up spectrum taken with Keck showed the spectral type to be M9; from the previous USNO and DENIS imaging, a high proper motion of $1.516 \pm 0.012''$ per year to the SE was determined; from its

magnitudes and spectral type, a distance of 4.1 ± 0.6 pc was determined; its galactic UVW velocities are typical for the thin disk (Delfosse et al. 2001). Deacon & Hambly (2001) obtained its trigonometric parallax from archived plates to be 192 ± 37 mas, i.e. a distance of $5.2^{+1.2}_{-0.8}$ pc, still consistent with the photometric distance estimate from Delfosse et al. (2001) for a single or binary star.

Because of its small distance and intrinsically faint magnitude, such an object is a promising target for direct imaging detection of sub-stellar companions, both brown dwarfs and giant planets. Due to the problem of dynamic range, namely that sub-stellar objects are too faint and too close to their primary stars, it is difficult to directly detect such objects in orbit around a normal star. One way around this problem is to observe intrinsically faint objects like so-called ultra-cool dwarfs (i.e. dwarfs with spectral type M6 or later, including brown dwarfs). If they are also very nearby, one can probe

Correspondence to: rne@mpe.mpg.de

^{*} Data presented here were obtained in ESO programs 66.D-0135(A), 66.C-0138(B), 67.C-0325(A), 67.C-0514(A), and 67.C-0354(A) on La Silla and Cerro Paranal, as well as with the 2.2m telescope of the University of Hawai'i Institute for Astronomy on Mauna Kea.

separations which are of the order of the semi-major axes of the giant planets of our own solar system.

So far, only a few brown dwarf companion candidates detected directly in orbit around normal stars are confirmed by both spectroscopy and proper motion, the first of which was Gl 229 B (Nakajima et al. 1995, Oppenheimer et al. 1995), and the youngest of which is TWA-5 B (Lowrance et al. 1999, Neuhäuser et al. 2000). The first four confirmed brown dwarf companions¹ all orbit M-type dwarf stars. Only the fifth such companion, HR 7329 B, orbits an A0-type star (Lowrance et al. 2000, Guenther et al. 2001). This may indicate that nature prefers binaries with not too different masses, another motivation to search for sub-stellar companions around M-type stars; however, it could also be a selection effect, because companions with very different masses and magnitudes are much more difficult to detect.

In Sect. 2, we show our first two deep infrared (IR) imaging observation of Denis 1048–39. Follow-up H- and K-band spectroscopy is presented in Sect. 3. Two more follow-up images of Denis 1048–39 are used to derive the proper motion of the companion candidate in Sect. 4 and the parallax of Denis 1048–39 itself in Sect. 5. Then, in Sect. 6, we search for closer companions with the speckle camera SHARP-I. We discuss and summarize our results in Sect. 7.

2. The first two deep imaging observations

Deep imaging with high dynamic range can best be done in the near IR, because of low magnitude difference between the primary and possible sub-stellar companions, good sky conditions like seeing (compared to, e.g., the optical), and high sensitivity of near IR detectors (compared to, e.g., the thermal IR).

We observed Denis 1048–39 for the first time with the Son of Isaac (Sofi²) at the 3.5m New Technology Telescope (NTT) of the European Southern Observatory (ESO) on La Silla, Chile, during the morning twilight of 8 December 2000 (at airmass 1.0), while the targets of that program (companions to nearby neutron stars) were not observable anymore. We used the small Sofi field with $0.147''$ per pixel for better angular resolution and obtained 30×30 sec integrations. Darks, flats, and standards were observed in the same night with the same set-up and data reduction was done with *eclipse*³. We measured a FWHM of $0.51''$ in our Denis 1048–39 images, see Table 1 for the observations log.

Using the faint HST standard stars S427-D and S341-D, we obtain the as yet unknown H-band magnitude for Denis 1048–39, namely $H=9.55 \pm 0.16$ mag. This value is the mean of three measurements taken with Sofi (see Table 1 for the observation log), which individually show relatively large errors (~ 0.15 mag each) and a large scatter (9.67, 9.37, and 9.60 mag, still consistent with being constant), because we




Fig. 1. NTT/Sofi H-band image of Denis 1048–39 with surrounding field, taken on 8 Dec 2000. The faint companion candidate is clearly seen $\sim 6''$ NNW of the primary. The other marked stars are used as background comparison stars for the measurement of proper and parallactic motion.

Table 1. Observations log

Telescope	instrument	obs. date	filter	FWHM
NTT ¹	Sofi	8 Dec 2000	H	$0.51''$
2.2m ²	QUIRC	15 Mar 2001	H	$1.20''$
NTT ¹	Sofi	6 Apr 2001	H ⁴	$1.36''$
VLT ³	ISAAC	21 Apr 2001	H & K ⁵	$0.40''$
NTT ¹	Sofi	9 Jun 2001	H	$0.70''$
NTT ¹	SHARP-I	2 Jul 2001	K	$0.34''$

Remarks: (1) La Silla. (2) Mauna Kea. (3) Cerro Paranal. (4) Companion candidate not detected due to insufficient dynamic range. (5) Companion candidate not detected in acquisition image because of very short integration time (1.77 sec).

observed only few photometric standard stars (aiming at pure detections of companion candidates rather than precise photometry), so that the mean of the three measurements also has a relatively large error, even though the star is quite bright. Our H-band magnitude is roughly consistent with its previously known BRIJK colors and its spectral type M9, as well as a distance of roughly 4 to 5 pc ($M_H \simeq 10.9$ mag for few Gyr old M9 dwarf stars according to Kirkpatrick & McCarthy 1994).

We detected a faint object near Denis 1048–39 (Fig. 1) with $H=15.9 \pm 0.5$ mag, i.e. 6.4 ± 0.5 mag fainter than Denis 1048–39. In December 2000, the companion candidate was located $5.766 \pm 0.014''$ NNW of Denis 1048–39 (see Table 2).

If the companion candidate would be bound, it should be a sub-stellar object. Whether the candidate indeed is bound, can be checked best with a 2nd epoch image to measure its proper motion.

¹ in addition to Gl 229 B and TWA-5 B mentioned above, there are G 196-3 B (Rebolo et al. 1998) and GD 570 D (Burgasser et al. 2000)

² see www.la.eso.org/lasilla/Telescopes/NEWNTT/

³ see www.eso.org/projects/aot/eclipse/

We obtained our 2nd epoch image with the QUick In-fraRed Camera (QUIRC⁴), a $1k \times 1k$ Hawaii array, at the 2.2m (88 inch) telescope of the University of Hawai'i Institute of Astronomy on Mauna Kea in the nights of March 13/14 and 14/15. The images of both nights were flat fielded, sky subtracted, shifted, and co-added for the two nights separately with MIDAS⁵. Because the seeing was better in the second of those two nights, we use only the data of that night (174 co-added images with 5 sec integration time each). Because of the relatively large airmass (~ 2) of Denis 1048–39 for our Mauna Kea observation, the FWHM is not very good anyway (Table 1).

The companion candidate is marginally detected in the QUIRC image. Pixel scale ($0.216''$ per pixel) and orientation (north is up, but tilted by 5.5° to the left) of the detector QUIRC were measured with our image of the binary star GJ 490 obtained in the same night with QUIRC (compared to its 2MASS image⁶). The separation between primary object and companion candidate ($6.0 \pm 0.3''$) in this 2nd epoch image is not inconsistent with the 1st epoch image (Fig. 1), so that it could be a co-moving pair, even though the significance is low because of the large FWHM in the 2nd epoch image.

3. Spectroscopic follow-up observation

H- and K-band spectra were taken with the Infrared Spectrograph and Array Camera (ISAAC) at the ESO 8.2m telescope Antu, Unit Telescope No. 1 (UT1) of the Very Large Telescope (VLT) on Cerro Paranal. Both spectra were obtained on 21 April 2001 in service mode and consist of 40 co-added 60 sec exposures each through a $1''$ slit. The slit was aligned neither along the position angle of the primary and its companion candidate, nor perpendicular to that angle, but in between those two angles, so that we could detect the signal from the companion candidate in the wing of the PSF of the primary: If the slit would have been aligned along the position angle, the primary would have been too bright, and if we would have aligned the slit perpendicular to that angle, the signal of the faint companion candidate would have been on top of the bright signal from the primary in the collapsed spectral profile. An angle in between the two extremes is the best compromise according to our experience (e.g. Neuhauser et al. 2001). Darks, flats, and arcs were taken in the same night.

The target aquisition was done in the following way: First, a short aquisition image was taken; then, the slit was aligned in north-south direction (as explained above) and centered on the primary; then, a small offset to the west was applied as given by the separation of the companion candidate observed in Dec 2000. Such an aquisition sequence with the VLT is usually very precise and accurate. If the companion candidate would indeed be bound, it should have the same proper and parallactic motion as the primary.

After standard data reduction (dark subtraction, normalization, flat fielding, sky subtraction, wavelength calibration,

Table 2. Separations of the companion candidate

Obs. date	$\Delta\alpha$	$\Delta\delta$	separation
8 Dec 2000	2.510(24)'' W	5.191(13)'' N	5.766(14)''
9 Jun 2001	1.517(25)'' W	5.432(21)'' N	5.640(18)''

and co-adding the spectra), we have searched for the faint signal from the companion candidate in the wing of the bright primary's PSF. The separation between the peak of the primary in the slit and the peak of the companion candidate should be 35 pixel (i.e. $5.2''$), the separation projected onto the slit orientation. However, we could not find the faint companion candidate's signature. This can be taken as first evidence for the companion candidate to be an unrelated, namely non-moving, background object: If it would have been a co-moving companion, it should have been in the slit⁷.

The H- and K-band spectra, which we obtained for the Denis 1048–39 primary, even though located slightly outside the slit, are typical for an M9 dwarf as far as the general shape of the continuum and the presence and strength of absorption lines (e.g. Na and CO) are concerned.

4. Proper motion of the companion candidate

Because the image obtained at Mauna Kea has a large FWHM, we had to take another image to measure the proper motion with higher significance. We tried to obtain such an image of Denis 1048–39 and its companion candidate on 6 April 2001 with SofI at the NTT (at airmass 1.2), while the targets of that program (in the ρ Ophiuchi dark cloud) were not visible at good airmass. However, due to unfavorable seeing of $\sim 1.4''$, the companion candidate is not detected in those images (30×30 sec each in H with the small SofI field, Table 1). The Denis 1048–39 primary itself and several other background comparison stars are well detected in this image, which we will use below for the determination of the parallax of Denis 1048–39.

Then, we took another image on 9 June 2001 (at airmass 1.4), again with SofI at the NTT, while the targets of that program (dark clouds) were too low (again 30×30 sec in H with the small SofI field, Table 1). This image is useful to measure the proper motion of the companion candidate (by comparison with the 1st image, Table 2) and also to determine the parallax of the Denis 1048–39 primary – together with the previous two SofI images obtained in $\sim 1/4$ year intervalls, see Sect. 5.

The two separations measured between the primary and the companion candidate as obtained from the 1st and 4th image (both with SofI) are listed in Table 2. While the errors in $\Delta\alpha$ and $\Delta\delta$ given in Table 2 take into account an uncertainty in the orientation of the detector (to the north) of $\pm 0.01''$, the error in separation does not include that error, because the separation can be measured independant of the orientation.

⁷ The same argument, the other way around, was used by Lowrance et al. (2000) for the HR 7329 B companion candidate: Because it was in the slit after blind offset, it is probably a co-moving companion, which was later confirmed by Guenther et al. (2001).

⁴ see <http://www.ifa.hawaii.edu/88inch/>

⁵ see www.eso.org/projects/esomidas/

⁶ see www.ipac.caltech.edu/2mass/

All errors given in Table 2 include an uncertainty in the pixel scale of 1 %.

The separation between the primary and the companion candidate has changed significantly from Dec 2000 and June 2001. This cannot be explained by orbital motion after half a year. Hence, the two objects do not form a common proper motion pair. The companion candidate is clearly a background object.

Assuming that the companion candidate is a non-moving background object, the observed motion of Denis 1048–39 itself (to the SW) relative to this background object from Dec 2000 to June 2001 (Table 2) is consistent with its known proper motion (Delfosse et al. 2001), taking into account some parallactic motion of Denis 1048–39 (see below).

5. The parallax of Denis 1048–39

For determining the parallax of Denis 1048–39, we measured the separations in α and δ between Denis 1048–39 and ten comparison stars (marked in Fig. 1) including all separations between comparison stars, in the same way for all three images taken with SofI. First, we checked that the separations between the comparison stars did not change. The precision in the Gaussian centering of the comparison stars and the target ranges from 0.16 pixel (24 mas) in the best image (Dec 2000) to 0.33 pixel (49 mas) in the April 2001 image. Hence, the precision in one measurement of a parallactic displacement (i.e. the difference between two stars from one image to another one) ranges from 0.5 to 0.7 pixel, i.e. 74 to 103 mas. When we average the relative motion (from Dec 2000 to April 2001) between the target and several comparison stars, the precision of the target motion is again better, namely ~ 33 mas in any one direction.

The (averaged) motion of Denis 1048–39 with respect to the comparison stars from Dec 2000 to Apr 2001 is $0.750 \pm 0.043''$ to the west and $0.321 \pm 0.101''$ to the south. From this motion, we have to subtract the proper motion (known from Delfosse et al. 2001, and Deacon & Hambly 2001 to be $\mu\alpha = -1.147 \pm 0.008''$ and $\mu\delta = -0.992 \pm 0.009''$). The remaining parallactic motion from Dec to April (0.326 yrs) is $0.376 \pm 0.043''$ to the west and $0.002 \pm 0.101''$ to the north. Similarly, the parallactic motion from April to June (0.175 yrs) is $0.151 \pm 0.062''$ to the west and $0.141 \pm 0.023''$ to the north.

One can estimate the parallactic motion for each position in the sky and any set of observing dates. For the Denis 1048–39 position and our observing dates, we can estimate the parallactic motion in right ascension and declination for a range of possible distances, e.g. from 4 to 6 pc. Those estimates have to be compared with the observed motion to derive the true distance.

In Fig. 2, we show the *difference* between expected and observed parallactic motion (in α , δ , and total) for a range in distance. The total motion (Dec 2000 to June 2001) is best consistent with a distance of ~ 4.6 pc. Given our astrometric precision of ~ 33 mas (see above), the error is ± 0.3 pc, as seen in Fig. 2. We have also measured the parallactic motion from the Dec to April and then from the April to June,




Fig. 2. Determining the parallax of Denis 1048–39. We plot the difference (in mas) between observed and expected parallactic motion (from Dec 2000 to June 2001) for a range in distance: Dashed line for motion in right ascension α , dotted line for declination δ , and full line for total motion. The difference in total motion is smallest at ~ 4.6 pc, which is therefore the observed distance towards Denis 1048–39. Given our astrometric precision of 33 mas (see text), the error is ± 0.3 pc, indicated by the box.

both are consistent with each other and with the value given above for the Dec 2000 to June 2001 total motion. Also, the individual motion (Dec 2000 to June 2001) in both α and δ are consistent with 4.6 ± 0.3 pc within the errors. This new distance is more precise than previously known and also consistent with the previous estimates.

6. Follow-up speckle imaging of Denis 1048–39

In the evening twilight of 2 July 2000, we observed Denis 1048–39 using SHARP-I (System for High Angular Resolution Pictures, Hofmann et al. 1992) at the NTT. We have obtained a total of 1500 images of 0.5 sec exposure each in the K-band, spread into six parts of 250 images; from one such part to the next, we moved back and forth on two detector quadrants. According to the La Silla seeing monitor, the average seeing during the observations was $0.86''$. The FWHM in the final co-added image is $0.34''$ (using the nominal SHARP pixel scale of $0.049''$). The data were corrected for bad pixels followed by a sky image subtraction and the application of a flat-field. For each band we then co-added the 256×256 pixel frames using the brightest pixel as shift-and-add reference (Christou 1991).

We reduced the SHARP-I speckle data following Köhler et al. (2000) to compute the limiting magnitude for undetected but detectable close companion candidates from the visibilities, the Knox-Thompson phase, and the bispectrum, the first of which clearly gives the lowest limits. The limits for the brightness ratio K_1/K_2 between primary (K_1) and any companion candidates (K_2) for the six data cubes range between 0.03 and 0.12. When averaging all six cubes, the limit to K_1/K_2 is 0.01 for any companion candidate outside of 3 pixels. We present the achieved dynamic range for the speckle




Fig. 3. Dynamic ranges achieved. We plot the log of the flux ratio between the 3σ background noise level and the peak intensity of Denis 1048–39 (and, on the right hand side, the magnitude difference) versus the separation to the primary’s photocenter (and, on the top axis, the projected physical separation at 4.6 pc), for the three detectors used (from top to bottom): QUIRC at the Mauna Kea 2.2m, Sofi at the La Silla NTT (both in H) as well as SHARP-I at the NTT (in K); for SHARP-I, we plot the curve obtained from the simple shift and add image (from 0 to $3''$) and the visibility curve (vis) calculated following the Köhler et al. (2000) procedure (for the inner $2''$), the latter of which has better dynamic range for the inner $\sim 1.1''$. The detected companion candidate found to be an unrelated background object is indicated as cross ($\sim 2\sigma$ detection with QUIRC). The dotted lines show the expected flux ratios for 10 and $25 M_{\text{jup}}$ mass companions at 1 and 2 Gyrs around the primary (in the H-band) calculated from Burrows et al. (1997) models. Non-detections of bound companions outside of $\sim 0.25''$ (or $2''$) exclude any stellar (or any brown dwarf, respectively) companions outside ~ 1 (or 9) AU (at 2 Gyrs). From the SHARP-I visibility curve, we can exclude companions down to a flux ratio of 1:100 ($\sim 40 M_{\text{jup}}$ at 2 Gyrs) outside of $0.15''$ (0.7 AU).

data in Fig. 3, both for the visibility computed following Köhler et al. (2000) as well as the dynamic range achievable by simple shift and add. From the visibility curve, we can exclude any companion candidate outside of $0.15''$ (i.e. 0.7 AU at 4.6 pc) down to a limiting magnitude of ~ 13.6 mag in K (with $K = 8.58 \pm 0.05$ mag for the primary, Delfosse et al. 2001), which corresponds to a $\sim 40 M_{\text{jup}}$ (Burrows et al. 1997).

The previous companion candidate discussed above is not detected, because we placed the primary only in the two lower, i.e. northern SHARP-I quadrants, because they have better pixel and flat field characteristics; each quadrant has a field-of-view of $\sim 6.2'' \times 6.2''$. With the SHARP-I observations, we concentrated on detecting even closer com-

panion candidates, within $3''$; there was no reason to image the previously rejected companion candidate again. No additional companion candidates are detected in the $12.5'' \times 12.5''$ SHARP-I field-of-view. When comparing the PSF of Denis 1048–39 with the PSF reference star PPM 288121 (a known single star observed immediately before and after the prime target), we find no evidence for any elongation of the Denis 1048–39 PSF.

In Fig. 3, we compare the dynamic range achieved in our Sofi, QUIRC, and SHARP images. The flux ratio is determined from our actual images of Denis 1048–39 as the 3σ background noise level (on 49 pixels) divided by the peak intensity. We compare the observed dynamic ranges with expected flux ratios for possible companions of different masses (calculated following Burrows et al. 1997), for an age of 1 and 2 Gyrs for the primary at 4.6 pc distance (with $B.C.H = 2.6$ mag as for Gl 229 B, Leggett et al. 1999). Hence, for an age of 2 Gyrs, we should have detected all companions with mass above $\sim 10 M_{\text{jup}}$ outside of $5''$ (23 AU) and all companions with mass $\geq 25 M_{\text{jup}}$ at $\geq 3''$ separation (14 AU), all detectable in the Sofi images. The MPE speckle camera SHARP-I clearly gives the best dynamic range. In the SHARP images, we should have detected all $\sim 10 M_{\text{jup}}$ companions between 2 and $3''$ separation (9 to 14 AU) and companions with masses $\geq 25 M_{\text{jup}}$ at $\geq 1''$ separation (5 AU). If Denis 1048–39 is a $\sim 0.09 M_{\odot}$ mass star (see below), any companion which is ≥ 1 mag fainter would be sub-stellar (Chabrier et al. 2000). Outside of $0.15''$ (0.7 AU), we should have detected all companions with magnitude difference of $\Delta K \geq 5$ mag with SHARP-I.

7. Discussion and Summary

A faint, possibly sub-stellar companion candidate $6''$ NNW of Denis 1048–39 turned out to be a background object. Because several other groups are actively searching for such sub-stellar companions around nearby stars, we think that it is important also to report such a negative result. Because the primary object itself is almost sub-stellar, any real companion which is just a few mag fainter, would be sub-stellar. Because of the high dynamic range achieved with the speckle camera SHARP-I, we can exclude stellar companions at separations above $0.25''$, which is only 1 AU at 4.6 pc distance, and we can also exclude brown dwarf companions at $\geq 2''$ (9 AU), all for an age of 2 Gyrs.

Three epoch imaging led to the determination of the parallax of Denis 1048–39 corresponding to a distance of 4.6 ± 0.3 pc. This distance estimate is consistent with both the photometric distance of 4.1 ± 0.6 pc given in Delfosse et al. (2001) in case that the object is a single main sequence M9-dwarf as well as with the trigonometric parallax measurement (from archived plates) by Deacon & Hambly (2001). At this distance, its apparent magnitudes (assuming no interstellar reddening) the following absolute magnitudes, rarely available for an M9-dwarf: $M_B = 20.69 \pm 0.33$, $M_R = 17.39 \pm 0.24$, $M_I = 14.36 \pm 0.14$, $M_J = 11.28 \pm 0.14$, $M_H = 11.24 \pm 0.34$, and $M_K = 10.27 \pm 0.14$ mag.

At the given distance, its colors and spectral type place the object into the H-R diagram, so that we can compare its position with theoretical models (Burrows et al. 1997, Chabrier et al. 2000). The object is a very low-mass star with a mass of ~ 0.075 to $\sim 0.09 M_{\odot}$ and an age of ~ 1 to 2 Gyrs. There is no star known that is both more nearby and cooler than Denis 1048–39. At a distance of ~ 4.6 pc, it is the 40th nearest star.⁸

Speckle imaging with SHARP has revealed no additional closer companion candidates, so that it seems likely that Denis 1048–39 is a single star. However, an even lower mass or even closer companion cannot be excluded.

Our direct imaging observations of the companion candidate at $\sim 6''$ separation with an H-band magnitude difference of ~ 6.4 mag shows that such faint possibly sub-stellar companions can be detected directly around nearby ultra-cool dwarfs, in this case at a projected physical separation of only ~ 27 AU, if it were bound. In the near future, this target and other similar nearby ultra-cool dwarfs can be observed with the repaired NICMOS at the HST and CONICA-NAOS at the VLT, which provides an IR wavefront sensor. Then, we can investigate the multiplicity of such objects with even higher dynamic range and higher sensitivity.

Acknowledgements. We would like to thank the ESO User Support Group for assistance, the ISAAC team for the VLT service mode observations, and the NTT team with O. Hainaut, L. Vanci, G. Martin, and J. Miranda for support during the SofI observations. We are grateful to Klaus Bickert and Rainer Schödel for their help with the Sharp run. We also thank Eduardo Martín for introducing Denis 1048–39 to us shortly after its publication on astro-ph and in a press release in November 2000. RN and WB would like to thank the Institute for Astronomy of the University of Hawai'i for hospitality, where they have done part of this research, as well as for allocation of several observing nights with its 2.2m telescope on Mauna Kea. We also thank Susan Parker and Andrew Pickles for their help with those observations. RN wishes to acknowledge financial support from the Bundesministerium für Bildung und Forschung through the Deutsche Zentrum für Luft- und Raumfahrt e.V. (DLR) under grant number 50 OR 0003. The authors wish to extend special thanks to those Hawai'ians, on whose sacred mountain we worked (without having asked).

References

- Burgasser A.J., Kirkpatrick J.D., Cutri R.M. et al. 2000, ApJ 531, L57
- Burrows A., Marley M., Hubbard W. et al. 1997, ApJ 491, 856
- Chabrier G., Baraffe I., Allard F., Hauschildt P., 2000, ApJ 542, 464
- Christou J.C., 1991, Experimental Astr. 2, 27
- Deacon N.R. and Hambly N.C., 2001, A&A 380, 148
- Delfosse X., Forveille T., Martín E.L., et al., 2001, A&A 366, L13
- Epchtein N., de Batz B., Capoani L., 1997, The ESO Messenger 87, 27
- Guenther E.W., Neuhauser R., Huélamo N., Brandner W., Alves J., 2001, A&A 365, 514
- Hofmann R., Blietz M., Duhoux Ph, Eckart A., Krabbe A., Rotaciuc V., 1992, SHARP and FAST: NIR Speckle and Spectroscopy at the MPE. In: *Progress in Telescope and Instrumentation Technologies*, Ulrich M.-H. (Ed.), ESO Conference and Workshop Proceedings No. 42, 617
- Kirkpatrick J.D. and McCarthy D.W., 1994, AJ 107, 333
- Köhler R., Kunkel M., Leinert C., Zinnecker H., 2000, A&A 356, 541
- Lowrance P.J., McCarthy C., Becklin E.E. et al., 1999, ApJ 512, L69
- Lowrance P.J., Schneider G., Kirkpatrick J.D. et al., 2000, ApJ 541, L390
- Nakajima T., Oppenheimer B.R., Kulkarni S.R. et al., 1995, Nat 378, 463
- Neuhäuser R., Brandner W., Eckart A., et al., 2000, A&A 354, L9
- Neuhäuser R., Guenther E.W., Brandner W., et al., 2001, *Direct imaging search for planetary companions next to young nearby stars*. In: From Darkness to Light, ed. by Montmerle T. & André P., ASP Conf. Series 243, 723
- Oppenheimer B.R., Kulkarni S.R., Matthews K., van Kerkwijk M.H., 1995, Sci 270, 1478
- Rebolo R., Zapatero-Osorio M.R., Madrugá S. et al., 1998, Sci 282, 1309

⁸ see <http://www.chara.gsu.edu/RECONS/TOP100.htm>